



NYSERDA

New York State Greenhouse Gas Inventory: 1990 – 2014

Final Report

NYSERDA's Promise to New Yorkers:

NYSERDA provides resources, expertise, and objective information so New Yorkers can make confident, informed energy decisions.

Mission Statement:

Advance innovative energy solutions in ways that improve New York's economy and environment.

Vision Statement:

Serve as a catalyst – advancing energy innovation, technology, and investment; transforming New York's economy; and empowering people to choose clean and efficient energy as part of their everyday lives.

NYSERDA Record of Revision

Document Title
New York State Greenhouse Gas Inventory: 1990-2014 December 2016 Revised February 2017

Revision Date	Description of Changes	Revision on Page(s)
Add Date	Original Issue	December 2016
2/2/2017	Section S.3.1: Added sentence to end of paragraph 2, and added corresponding footnote	Page S-4
2/2/2017	Section 3.3 Results: replaced paragraph text	Page 25

New York State Greenhouse Gas Inventory: 1990–2014

Final Report

Prepared by:

New York State Research and Development Authority

Albany, NY

December 2016
Revised February 2017

Table of Contents

NYSERDA Record of Revision	i
List of Figures	v
List of Tables	v
Acronyms and Abbreviations	vi
Summary	S-1
1 Residential, Commercial/Institutional, and Industrial Fuel Combustion.....	1
1.1 Overview	1
1.2 Emissions Inventory Data and Methodology.....	1
1.3 Results	3
1.3.1 Residential Sector	3
1.3.2 Commercial/Institutional Sector	6
1.3.3 Industrial Sector	8
2 Industrial Non-Fuel Combustion Processes	11
2.1 Overview	11
2.1.1 Sources of Carbon Dioxide (CO ₂) Emissions	11
2.1.2 Sources of Methane (CH ₄) Emissions	12
2.1.3 Sources of Perfluorocarbon (PFC) Emissions	13
2.1.4 Sources of Hydrofluorocarbon (HFC) Emissions	14
2.1.5 Other Sources of Non-Fuel Combustion Emissions	14
2.2 Emissions Inventory Data and Methodology.....	14
2.2.1 Sources of CO ₂ Emissions.....	15
2.2.2 Sources of CH ₄ Emissions.....	16
2.2.3 Sources of PFC Emissions	17
2.2.4 Sources of HFC Emissions	17
2.3 Results	17
2.3.1 Sources of CO ₂ Emissions.....	19
2.3.2 Sources of CH ₄ Emissions.....	20
2.3.3 Sources of PFC Emissions	20
2.3.4 Sources of HFC Emissions	20
3 Transportation Energy Use	21
3.1 Overview	21
3.1.1 On-Road Vehicle Fuels	21
3.1.2 Off-Road Vehicle Fuels	22

3.1.3	Lubricants	22
3.2	Emissions Inventory Data and Methodology.....	23
3.2.1	On-Road Vehicle Fuels	23
3.2.2	Off-Road Vehicle Fuels	24
3.2.3	Lubricants	25
3.3	Results	25
4	Power Supply and Delivery	27
4.1	Overview	27
4.1.1	Fuel Combustion	27
4.1.2	Electricity Distribution	27
4.1.3	Municipal Waste Combustion.....	28
4.2	Emissions Inventory Data and Methodology.....	28
4.2.1	Fuel Combustion	28
4.2.2	Electricity Distribution	29
4.2.3	Municipal Waste Combustion.....	29
4.3	Results	30
5	Agriculture.....	35
5.1	Overview	35
5.1.1	Livestock	35
5.1.2	Agricultural Soils.....	36
5.2	Emissions Inventory Data and Methodology.....	36
5.2.1	Livestock	37
5.2.2	Agricultural Soils.....	38
5.3	Results	38
6	Waste Management	40
6.1	Overview	40
6.1.1	Landfills	40
6.1.2	Municipal Wastewater Management.....	40
6.2	Emissions Inventory Data and Methodology.....	41
6.2.1	Landfills	41
6.2.2	Wastewater Management	41
6.3	Results	42
Appendix A	A-1	

List of Figures

Figure 1-1. Residential Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014	4
Figure 1-2. Residential Sector Heating Demand and GHG Emissions, 1990–2014	6
Figure 1-3. Commercial/Institutional Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014	7
Figure 1-4. Industrial Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014	9
Figure 2-1. GHG Emissions from Industrial Non-Fuel Combustion Processes by Source, 1990–2014	18
Figure 3-1. Transportation GHG Emissions by Fuel, 1990–2014	26
Figure 4-1. Proportion of New York State Electricity Generation and Net Imports by Source Category, 2014.....	30
Figure 4-2. Proportion of GHG Emissions from Power Supply and Delivery by Source Category, 2014.....	31
Figure 4-3. New York State Electricity Generation and Net Imports of Electricity by Source Category, 1990–2014.....	32
Figure 4-4. Primary Fossil Fuel Energy Use at New York State Power Stations by Fuel Type ..	33
Figure 4-5. GHG Emissions from Power Supply and Delivery by Source Category, 1990–2014	34
Figure 5-1. GHG Emissions from Agriculture, 1990–2014	39
Figure 6-1. GHG Emissions from Waste Management, 1990–2014	42

List of Tables

Table 1-1. 2014 Fuel Types for the Industrial Sector	2
Table 1-2. Residential Sector Emissions Inventory, 1990–2014 (MMtCO ₂ e).....	4
Table 1-3. Residential Sector Proportions of Total Emissions by Fuel Type, 1990–2014 (%).....	5
Table 1-4. Commercial/Institutional Sector Emissions Inventory, 1990–2014 (MMtCO ₂ e).....	7
Table 1-5. Commercial/Institutional Sector Proportions of Total Emissions by Fuel Type (%)	8
Table 1-6. Industrial Sector Emissions Inventory, 1990–2014 (MMtCO ₂ e).....	9
Table 1-7. Industrial Sector Proportions of Total Emissions by Fuel Type (%)	10
Table 2-1. Emissions Inventory for Industrial Non-Fuel Combustion Processes, 1990–2014 (MMtCO ₂ e)	19
Table 3-1. New York State VMT by Vehicle Type (million miles)	24
Table 3-2. Transportation Sector Emissions Inventory, 1990–2014 (MMtCO ₂ e).....	26
Table 4-1. Electric Generation (GWh) and Annual Average Growth Rates by Fuel Type	32
Table 4-2. GHG Emissions from Power Supply and Delivery by Source Category, 1990–2014 (MMtCO ₂ e)	34
Table 5-1. GHG Emissions from Agriculture, 1990–2014 (MMtCO ₂ e).....	39
Table 6-1. SIT Key Default Values for Municipal Wastewater Treatment.....	41
Table 6-2. GHG Emissions from Waste Management (MMtCO ₂ e), 1990–2014	42

Acronyms and Abbreviations

AEO	Annual Energy Outlook, published by the EIA
Btu	British thermal units
CAFE	Corporate average fuel economy
CH ₄	Methane
CNG	Compressed natural gas
CO ₂	Carbon dioxide
CO ₂ e	Carbon dioxide equivalent
CO ₂ FFC	Carbon dioxide from fossil fuel combustion
DOT	New York State Department of Transportation
EIA	Energy Information Administration
EPA	Environmental Protection Agency
FHWA	Federal Highways Administration
GSP	Gross State Product
GDP	Gross Domestic Product
GHG	Greenhouse Gas
Gwh	Gigawatt hours
HFC	Hydrofluorocarbons
HCFC	Hydrofluorochlorocarbons
LPG	Liquid petroleum gas
MMt	Million metric tons
MSW	municipal solid waste
N ₂ O	Nitrous oxide
NYISO	New York Independent System Operator
ODS	Ozone depleting substance
ORNL	Oak Ridge National Laboratory
PFC	Perfluorinated Chemicals
RCI	Residential, Commercial, Industrial
SEDS	State Energy Data System
SF ₆	Sulfur Hexafluoride
SIT	State Inventory Tool
T&D	Transmission and Distribution
VMT	Vehicle miles traveled

Summary

S.1 Introduction

Unequivocal warming of the Earth over the past century is documented by observations that include increases in global average temperatures, rapid melting of mountain glaciers and land ice sheets, and higher global average sea levels. In North America, extreme heat and drought events are becoming more frequent and prolonged. Although total precipitation is increasing only slightly, intense and damaging storms like Sandy and Irene are occurring more often. A changing climate affects human health, society, and the economy both directly and indirectly, through its effects on agriculture, sea level, fisheries, and other natural resources. The rate and extent of climate change depend on the amount of greenhouse gases (GHGs) present in, and delivered to, the atmosphere.

This report provides a detailed accounting of emissions in New York State from 1990–2014. The report identifies the emissions associated with different sectors and describes sector-specific calculation methodologies to provide greater detail and context on what sectors and sources are driving emissions. Accordingly, the chapters correspond to the different sectors, and are organized as follows:

- Chapter 1: Residential, Commercial/Institutional, and Industrial Fuel Combustion.¹
- Chapter 2: Industrial Non-Fuel Combustion Processes.²
- Chapter 3: Transportation Energy Use.
- Chapter 4: Power Supply and Delivery.
- Chapter 5: Agriculture.
- Chapter 6: Waste Management.

¹ Chapter 1 focuses on emissions in the Residential, Commercial, and Industrial sector, but also includes associated emissions from electricity generation that meets the demand for this sector. These electricity generation emissions are included in Chapter 1 for informational purposes only, as Chapter 4 separately identifies emissions from the electricity generation sector. While emissions from electricity generation appear in both chapters, they are not double counted.

² Chapter 2 provides estimates of emissions associated with natural gas T&D.

The report systematically looks at six primary GHG's: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF₆).

Emissions of these six GHGs are converted to and presented using a common metric, the CO₂ equivalent (CO₂e), so that the global warming potential of each of these different GHGs can be presented in equivalent terms.

S.2 New York State GHG Emissions: Sources and Trends

Table S-1 provides a summary of GHG emissions estimated for New York State by sector and gas in 2014, the most recent year for which historical data are available.

Table S-1. 2014 New York State Greenhouse Gas Inventory (MMtCO₂e)^a

	CO ₂	CH ₄	N ₂ O	PFC	HFC	SF ₆	Total	% of Total
Fuel Combustion^b								
(inc. Net Imports of Electricity)	179.69	0.41	0.88	-	-	-	180.98	83.12%
Electricity Generation	30.36	0.01	0.04	-	-	-	30.41	13.97%
Transportation	73.28	0.05	0.69	-	-	-	74.01	33.99%
Residential	35.19	0.25	0.06	-	-	-	35.50	16.31%
Commercial	21.93	0.07	0.02	-	-	-	22.03	10.12%
Industrial	10.98	0.02	0.04	-	-	-	11.04	5.07%
Net Imports of Electricity	7.96	0.01	0.03	-	-	-	7.99	3.67%
Fuel Combustion								
(exc. Net Imports of Electricity)	171.73	0.41	0.85	-	-	-	172.99	79.45%
Other Sources	3.94	19.74	2.44	0.38	10.03	0.22	36.74	16.88%
Power Supply & Delivery	2.69	-	0.05	-	-	0.22	2.96	1.36%
Electricity Transmission and Distribution	-	-	-	-	-	0.22	0.22	0.10%
Municipal Waste Combustion	2.69	-	0.05	-	-	-	2.74	1.26%
Agriculture, Forestry & Waste	-	17.56	2.39	-	-	-	19.95	9.16%
Agricultural Animals	-	3.56	-	-	-	-	3.56	1.63%
Agricultural Soil Management	-	-	1.46	-	-	-	1.46	0.67%
Landfills	-	11.63	-	-	-	-	11.63	5.34%
Manure Management	-	0.79	0.37	-	-	-	1.16	0.53%
Municipal Wastewater	-	1.58	0.55	-	-	-	2.13	0.98%
Industrial Processes & Manufacturing	1.24	2.18	-	0.38	10.03	-	13.83	6.35%
Aluminum Production	-	-	-	0.25	-	-	0.25	0.11%
Cement Production	0.38	-	-	-	-	-	0.38	0.18%
Iron & Steel Production	0.14	-	-	-	-	-	0.14	0.07%
Limestone Use	0.58	-	-	-	-	-	0.58	0.27%
Natural Gas Leakage ^b	-	2.18	-	-	-	-	2.18	1.00%
ODS Substitutes	-	-	-	-	10.03	-	10.03	4.61%
Semiconductor Manufacturing	-	-	-	0.13	-	-	0.13	0.06%
Soda Ash Use	0.13	-	-	-	-	-	0.13	0.06%
Total (inc. Net Imports of Electricity)	183.63	20.15	3.31	0.38	10.03	0.22	217.73	100%
% of Total								
(inc. Net Imports of Electricity)	84.34%	9.26%	1.52%	0.17%	4.61%	0.10%	100%	-
Total (exc. Net Imports of Electricity)	175.67	20.15	3.29	0.38	10.03	0.22	209.73	-

^a MMtCO₂e = million metric tons of carbon dioxide equivalent; CO₂ = carbon dioxide; CH₄ = methane; N₂O = nitrous oxide; PFC = perfluorocarbons; HFC = hydrofluorocarbons; SF₆ = sulfur hexafluoride

^b 2014 energy-related emissions, a category calculated as part of New York's State Energy Plan (SEP) that includes fuel combustion, net electricity imports, power supply & delivery, and natural gas leakage, totals 186.12 MMtCO₂e, or 85 percent of total emissions.

^c A recent study found that measured emissions of methane from natural gas systems are approximately 1.5 times greater than those published in the U.S. Environmental Protection Agency national GHG inventory. A commensurate scaling of this analysis would increase the emissions from natural gas leakage to 3.3 MMTonsCO₂e or one percent of total emissions. See Brandt, et al., 2014. "Methane Leaks from North American Natural Gas Systems." *Science* 343, February.

S.3 Emissions

S.3.1 Overview

The *New York State Greenhouse Gas Inventory: 1990–2014* is based on the U.S. Environmental Protection Agency (EPA) production-based methodology and tailored to estimate current emissions produced within the State’s boundaries. The inventory adopts the EPA protocols for identified emissions, as well as categorization for sources of emissions. Additionally, the inventory reflects emissions associated with the electricity sources used to meet all of New York State’s demands, corresponding to a consumption-based approach for the power supply and delivery sector (i.e., emissions associated with imported electricity).

This report illustrates the historical development of GHG emissions. Table S-2 shows the trend in New York State’s historical GHG emissions from 1990–2014. Emissions gradually increased from 1990, peaked in 2005, and then began to decline. In 2014 emissions were approximately eight percent lower than in 1990. This reduction stands in contrast to the seven percent national increase in total GHG emissions over the same time period. Energy-related emissions were thirteen percent lower in 2014 relative to 1990 levels.³

³ Energy-related emissions is a category calculated as part of New York’s State Energy Plan (SEP) that includes fuel combustion, net electricity imports, power supply & delivery, and natural gas leakage.

Table S-2. New York State GHG Emissions, 1990–2014 (MMtCO₂e)

Values for 1990–2014 are based on historical data. GHG = greenhouse gas; MMtCO₂e = million metric tons of carbon dioxide equivalent. Totals may not equal exact sum of subtotals shown in this table due to independent rounding.

Gas and Category	1990	1995	2000	2005	2010	2014
Carbon Dioxide	204.72	203.62	225.58	229.11	189.66	183.63
<i>Fuel Combustion</i>	<i>201.57</i>	<i>199.66</i>	<i>221.63</i>	<i>223.65</i>	<i>186.02</i>	<i>179.69</i>
Electricity Generation	62.76	51.10	55.47	53.38	37.20	30.36
Net Imports of Electricity	1.63	4.24	5.66	6.52	9.55	7.96
Transportation	57.18	60.72	71.77	81.14	73.58	73.28
Residential	33.76	34.37	39.40	39.22	31.42	35.19
Commercial	26.37	26.84	31.97	28.45	24.06	21.93
Industrial	19.86	22.40	17.37	14.94	10.22	10.98
<i>Other Sources</i>	<i>3.16</i>	<i>3.96</i>	<i>3.95</i>	<i>5.46</i>	<i>3.64</i>	<i>3.94</i>
Municipal Waste Combustion	1.23	1.90	2.00	3.52	2.30	2.69
Cement Production	0.67	0.76	0.80	0.79	0.51	0.38
Iron and Steel Production	0.83	0.83	0.75	0.60	0.24	0.14
Limestone Use	0.24	0.29	0.22	0.37	0.45	0.58
Soda Ash Use	0.20	0.19	0.18	0.17	0.14	0.13
Methane	23.52	24.57	23.95	22.29	20.22	20.15
<i>Fuel Combustion</i>	<i>0.68</i>	<i>0.75</i>	<i>1.04</i>	<i>0.73</i>	<i>0.39</i>	<i>0.41</i>
Electricity Generation	0.04	0.02	0.03	0.03	0.01	0.01
Net Imports of Electricity	0.00	0.00	0.00	0.00	0.01	0.01
Transportation	0.11	0.10	0.09	0.07	0.06	0.05
Residential	0.37	0.47	0.70	0.47	0.20	0.25
Commercial	0.11	0.13	0.18	0.14	0.08	0.07
Industrial	0.05	0.03	0.04	0.03	0.02	0.02
<i>Other Sources</i>	<i>22.84</i>	<i>23.82</i>	<i>22.91</i>	<i>21.55</i>	<i>19.83</i>	<i>19.74</i>
Agricultural Animals	3.37	3.20	3.30	3.17	3.36	3.56
Landfills	12.89	13.43	13.52	13.49	12.17	11.63
Manure Management	0.50	0.56	0.68	0.75	0.77	0.79
Municipal Wastewater	1.44	1.45	1.52	1.55	1.55	1.58
Natural Gas Leakage	4.63	5.17	3.88	2.59	1.99	2.18

Table S-2 continued

Gas and Category	1990	1995	2000	2005	2010	2014
Nitrous Oxide	5.93	6.26	6.37	5.10	3.89	3.31
<i>Fuel Combustion</i>	<i>3.51</i>	<i>4.11</i>	<i>4.26</i>	<i>2.89</i>	<i>1.49</i>	<i>0.88</i>
Electricity Generation	0.18	0.13	0.15	0.14	0.08	0.04
Net Imports of Electricity	0.01	0.01	0.02	0.02	0.03	0.03
Transportation	3.10	3.76	3.82	2.51	1.27	0.69
Residential	0.09	0.10	0.15	0.11	0.06	0.06
Commercial	0.05	0.05	0.06	0.05	0.03	0.02
Industrial	0.08	0.05	0.07	0.05	0.03	0.04
<i>Other Sources</i>	<i>2.41</i>	<i>2.15</i>	<i>2.11</i>	<i>2.21</i>	<i>2.39</i>	<i>2.44</i>
Agricultural Soil Management	1.59	1.32	1.26	1.30	1.51	1.46
Manure Management	0.31	0.29	0.27	0.30	0.30	0.37
Municipal Waste Combustion	0.04	0.06	0.05	0.08	0.04	0.05
Municipal Wastewater	0.47	0.49	0.52	0.52	0.54	0.55
Perfluorocarbons	0.44	0.41	0.47	0.37	0.42	0.38
Aluminum Production	0.38	0.31	0.33	0.27	0.32	0.25
Semiconductor Manufacturing	0.05	0.09	0.14	0.10	0.11	0.13
Hydrofluorocarbons						
ODS Substitutes	0.02	2.42	5.95	7.27	9.05	10.03
Sulfur Hexafluoride						
Electricity Transmission and Distribution	1.21	0.88	0.58	0.43	0.27	0.22
TOTAL	235.84	238.16	262.90	264.57	223.52	217.73
Totals	1990	1995	2000	2005	2010	2014
Fuel Combustion	205.76	204.52	226.93	227.27	187.90	180.98
Electricity Generation	62.99	51.25	55.65	53.55	37.29	30.41
Net Imported Electricity	1.63	4.26	5.69	6.55	9.59	7.99
Transportation	60.40	64.57	75.68	83.73	74.91	74.01
Residential	34.22	34.94	40.25	39.79	31.67	35.50
Commercial	26.53	27.01	32.21	28.64	24.17	22.03
Industrial	19.99	22.48	17.47	15.01	10.27	11.04
Other Sources	30.08	33.64	35.97	37.30	35.61	36.74
TOTAL	235.84	238.16	262.90	264.57	223.52	217.73
Energy-Related Emissions ^a	212.87	212.53	233.45	233.90	192.51	186.12


^a Energy-related emissions is a category calculated as part of New York's State Energy Plan (SEP) that includes fuel combustion, net electricity imports, power supply & delivery, and natural gas leakage.

S.3.2. Emissions Inventory

As shown in Table S-1, New York State accounted for approximately 218 million metric tons of carbon dioxide equivalent (MMtCO₂e) emissions in 2014, an average of 11 MtCO₂e for each State resident.⁴ At these levels, New York State's per capita GHG emissions were approximately half the U.S. average. The majority of the State's GHG emissions came from fuel combustion, which primarily represents the burning of fossil fuels (e.g., coal, natural gas, petroleum products) as an energy source to support various economic activities, including transportation, electric power generation, and heating and hot water needs for homes and businesses. Figure S-1 provides a breakdown of New York State's 2014 GHG emissions by gas. Even when considering the contributions of the six primary GHG emissions on a CO₂e basis, this figure shows that CO₂ contributes the majority (84 percent) of all GHG emissions in the State.

Figure S-1. 2014 Percentage of GHG Emissions by Gas and Source (Includes Net Imports of Electricity)

CO₂ = carbon dioxide; GHG = greenhouse gas.



⁴ Historic U.S. and New York State population data is drawn from the American Community Survey.


S.3.3 Fuel Combustion

S.3.3.1 Emissions by Economic Sector

The transportation sector accounted for approximately 41 percent of CO₂ emissions from fuel combustion in 2014 (shown in Figure S-2). The residential and commercial sectors were responsible for roughly 27 and 23 percent, respectively, after including an allocation of emissions from electricity generation. For both the residential and commercial sectors, emissions from on-site fuel combustion (including heating and hot water) were greater than the combined emissions associated with in-State and imported electricity generation. On-site fuel combustion from the industrial sector contributed approximately nine percent of the CO₂ fuel combustion emissions in the State.

Figure S-2. 2014 CO₂ Emissions from Fuel Combustion by End Use Sector (Includes Net Imports of Electricity)

CO₂ = carbon dioxide; GHG = greenhouse gas.




S.3.3.2 Emissions by Fuel

Total CO₂ emissions from fuel combustion were allocated to specific fuels, and then further distributed to the economic sector (transportation, electricity generation, residential, commercial, and industrial), providing a more detailed profile. Figure S-3 shows the fuels that contribute to the 2014 New York State CO₂ fuel combustion emissions. In 2014, natural gas accounted for 41 percent of CO₂ emissions. These emissions result primarily from natural gas combustion for electricity generation and on-site residential, commercial, and industrial use. An additional 30 percent of the CO₂ fuel combustion emissions result from the burning of gasoline by the transportation sector. Emissions contributions also remain across fossil fuels, due primarily to the burning of coal, distillate oil, aviation fuel, residual oil, diesel, and other petroleum sources, as well as imported electricity.

Figure S-3. 2014 CO₂ Emissions from Fuel Combustion by Fuel Type (Includes Net Imports of Electricity)

CO₂ = carbon dioxide; GHG = greenhouse gas.




S.3.3.3 Non-Fuel Combustion Emissions

Figure S-4 shows the sources of emissions from the non-fuel combustion category. In 2014, methane (which has a global warming potential 25 times greater than CO₂) accounted for the greatest portion of non-carbon dioxide CO₂e emissions. The majority of methane emissions are the result of activities other than fuel combustion, representing 9.1 percent of total statewide CO₂e emissions. The major sources of these methane emissions included landfills, natural gas system leakage, agricultural animals, and municipal wastewater facilities. Hydrofluorocarbon (HFC) emissions, resulting from use of substitutes for ozone-depleting substances (ODS), also represent a significant portion of non-fuel combustion emissions, at 4.6 percent of total statewide CO₂e emissions.

Figure S-4. 2014 Emissions from Non-Fuel Combustion Sources (Total Emissions Include Net Imports of Electricity)

CO₂ = carbon dioxide; GHG = greenhouse gas; ODS = ozone-depleting substance.




S.3.3.4 Emission Trends

Figure S-5 below illustrates several trends in New York State GHG emissions from 1990–2014. Of all in-State energy-related sources of emissions, the transportation sector emissions showed by far the greatest growth in New York State, with emissions increasing by nearly 23 percent. This is due to an increase in the consumption of gasoline and diesel fuels, as a result of an increase in vehicle miles traveled (VMT) in the State, and an increase in the consumption of jet fuel resulting from an increase in domestic air travel. Emissions from non-fuel combustion sources also increased, growing roughly 22 percent. This is primarily due to an increase in the use of HFCs as ODS substitutes.

In contrast, emissions from electricity generated in-State dropped 52 percent during this same period, acting as a major driver of the State’s decreasing GHG emissions. This drop is largely due to the significant decrease in the burning of coal and petroleum products in the electricity generation sector. Emissions from residential, commercial, and industrial buildings also decreased, showing a reduction of approximately 15 percent from 1990–2014. This reduction in emissions was primarily the result of a decrease in the use of coal and petroleum and an increase in the use of natural gas.

Figure S-5. GHG Emissions by Source Category, 1990–2014



S.3.3.5 GHG Emissions Intensity

New York State emits approximately 11 metric tons of CO₂e per capita (shown in Figure S-6), and its energy-related per capita emissions of 8.8 tons is the lowest of the 50 states. New York State also leads the nation in having the lowest GHG emissions per unit of economic output, averaging 0.17 kilograms (kg) of CO₂e of emissions per dollar gross state product (GSP), while the U.S. averaged 0.46 kg of CO₂e emissions per dollar gross domestic product (illustrated in Figure S-7).

Figure S-6. New York State and U.S. GHG Emissions Per Capita, 1990–2014





Figure S-7. New York State and U.S. Emissions Intensity



^a Gross state product (GSP) is the measure of state economic output, which is the sum of all value added by businesses in a state.

S.4 New York and U.S. Emission Comparisons

Figure S-8. 2014 GHG Emissions by Sector: New York State and U.S.

RCI = Residential, commercial/institutional, and industrial sector.




Figure S-8 shows the breakdown of emissions by major sector in New York State as compared to the United States. The principal sources of the State's GHG emissions in 2014 are the residential, commercial/institutional, and industrial sector; the transportation sector; and the power supply and delivery sector. Emissions for the residential, commercial, and industrial fuel use sectors are associated with the direct use of fuels (natural gas, petroleum, coal, and wood) to provide space heating, water heating, process heating, cooking, and other energy end-uses in the residential, commercial/institutional, and industrial sectors. This sector also accounts for GHG emissions from non-fuel sources in the industrial sector, such as CO₂ emissions from cement production, as well as emissions from the fossil fuel industry (e.g., natural gas leakage).

The transportation sector accounts for emissions associated with fuel consumption by all on-road and non-highway vehicles. Non-highway vehicles include jet aircraft, gasoline-fueled piston aircraft, railway locomotives, boats, and ships. The RCI sector includes emissions from non-highway agricultural and construction equipment. The power supply and delivery sector includes emissions associated with electricity generated within the State, imported electricity, emissions associated with municipal waste combustion (waste-to-energy facilities), and electricity transmission and distribution. The waste category includes emissions from landfills and wastewater. The U.S. agriculture emissions also include CH₄ and N₂O emissions from forest fires.

The residential, commercial, and industrial sector, the transportation sector, and the power supply and delivery sector account for 38, 34, and 19 percent of the State's GHG emissions, respectively. These sectors are also the three largest emitters in the U.S., but in a different order, with the power supply and delivery sector at 33 percent, the transportation sector at 30 percent, and the RCI sector at 28 percent. In New York State, emissions from waste and agriculture combine to account for the remaining nine percent of GHG emissions in 2014, while these two sectors account for nine percent of GHG emissions in the U.S. Overall, in 2014 New York State emissions accounted for three percent of U.S. total emissions.

S.5 A Closer Look at the Major Source of Emissions

The majority of the State's GHG emissions are associated with fuel combustion. The transportation sector accounts for the largest share of GHG emissions across the State, at 34 percent in 2014. Motor gasoline accounts for the majority of transportation GHG emissions; jet fuel contributes the second highest transportation GHG emissions; and diesel fuel ranks third among sources of transportation

emissions. Residual fuel, liquefied petroleum gas, and other transportation fuels account for the remaining transportation GHG emissions in 2014. Despite transportation emissions increasing from 1990–2014, emissions were highest in 2006 and have fallen by 13 percent from 2006 through 2014. The main drivers of both the 2006 peak and subsequent decline are consumption of motor gasoline, diesel fuel, and jet fuel.

Activities in the RCI⁵ fuel combustion sector produce GHG emissions when fuels are combusted to provide space heating, process heating, and other applications. Fuel combustion within the RCI sector accounts for 31 percent of the State’s GHG emissions in 2014, with a decrease in total emissions from 1990–2014. In 2014, the residential sector’s contribution toward the total RCI emissions from direct fuel use was 52 percent, while the commercial/institutional sector accounted for 32 percent and the industrial sector accounted for 16 percent.

In 2014, emissions from fuel combustion associated with the State’s electricity consumption are eight MMtCO₂e higher than those associated with in-State electricity production. The higher level for consumption-based emissions reflects GHG emissions associated with net imports of electricity.⁶ Electricity generation is dominated by natural gas and nuclear-powered units, with coal, oil, and hydro also important sources of historical generation in the State.

⁵ The industrial sector includes emissions associated with agricultural energy use and fuel used by the fossil fuel production industry.

⁶ Estimating the emissions associated with electricity use requires an understanding of the electricity sources (both in-State and out-of-State) used by utilities to meet consumer demand. For further details, see Chapter 4.

1 Residential, Commercial/Institutional, and Industrial Fuel Combustion

1.1 Overview

Activities in the Residential, Commercial/Institutional, and Industrial (RCI)⁷ sectors produce carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) emissions when fuels are combusted to provide space heating, water heating, process heating, cooking, and other energy end-uses. Carbon dioxide accounts for 99 percent of these emissions on a million metric tons (MMt) of CO₂ equivalent (CO₂e) basis in New York. In addition, since these sectors consume electricity, one can also attribute emissions associated with electricity generation to these sectors in proportion to their electricity use.⁸

1.2 Emissions Inventory Data and Methodology

Historical emissions estimations from direct fuel use were estimated using the EPA State Greenhouse Gas Inventory Tool (SIT) software and the methods provided in the Emission Inventory Improvement Program (EIIP) guidance document for RCI fuel combustion.⁹ The default historical fuel consumption data in SIT for New York, which came from the United States Department of Energy (DOE) Energy Information Administration's (EIA) *State Energy Data System* (SEDS), were updated with fuel consumption data for the Residential, Commercial/Institutional, and Industrial sectors from NYSERDA's *Patterns and Trends* report.¹⁰

⁷ The industrial sector includes emissions associated with agricultural energy use and fuel used by natural gas transmission and distribution (T&D) and oil and gas production industries.

⁸ Emissions associated with the electricity supply sector (presented in Chapter 4) have been allocated to each of the RCI sectors for comparison of those emissions to the fuel-consumption-based emissions presented in Chapter 1. Note that this comparison is provided for informational purposes and that emissions estimated for the electricity supply sector are not double-counted in the total emissions for the state. One could similarly allocate GHG emissions from natural gas T&D, other fuels production, and transport-related GHG sources to the RCI sectors based on their direct use of gas and other fuels, but we have not done so here due to the difficulty of ascribing these emissions to particular end-users. Estimates of emissions associated with the transportation sector are provided in Chapter 3, and estimates of emissions associated with natural gas T&D are provided in Chapter 2.

⁹ GHG emissions were calculated using the EPA's State Inventory Tool (SIT), with reference to: Emission Inventory Improvement Program (EIIP). 2006. "Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels." and "Chapter 2: Methods for Estimating Methane and Nitrous Oxide Emissions from Stationary Combustion." Volume VIII.

¹⁰ NYSERDA. 2016. "Patterns and Trends, New York State Energy Profiles: 2000-2014," (<https://www.nyserda.ny.gov/About/Publications/EA-Reports-and-Studies/Patterns-and-Trends>)

Note that the EIIP methods for the Industrial sector exclude the amount of stored carbon in products produced from fuels for non-energy uses from CO₂ emissions estimates. For example, the methods account for carbon stored in petrochemical feedstocks and in liquefied petroleum gases (LPG) and natural gas used as feedstocks by chemical manufacturing plants (i.e., not used as fuel), as well as carbon stored in asphalt and road oil produced from petroleum. The EIIP guidance document¹¹ explains the carbon storage assumptions for these products in detail. The primary tool to determine RCI combustion emissions is the Carbon Dioxide from Fossil Fuel State Inventory Tool Module (CO₂FFC SIT Module). In this module, fuel types are given carbon content values, combustion efficiencies,¹² and storage percentages. Table 1-1 shows fuel types for which the EIIP methods are applied in the SIT software to account for the Industrial Sector.

Table 1-1. 2014 Fuel Types for the Industrial Sector

Fuel Type
Coking Coal
Other Coal
Asphalt & Road Oil
Distillate Fuel
Feedstocks, Naphtha less than 401°F
Feedstocks, Other Oils greater than 401°F
Kerosene
Liquefied Petroleum Gas
Lubricants
Miscellaneous Petroleum Products
Petroleum Coke
Pentanes Plus
Residual Fuel
Special Naphthas
Waxes
Natural Gas

¹¹ Emissions Inventory Improvement Program. 2006. "Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels." Volume VIII.

¹² Combustion efficiency is defined as fuel specific percentage of carbon oxidized during combustion.

In addition to the fuel consumed directly by each of the RCI sectors, the proportion of each RCI sector's electricity sales to total electricity sales was used to allocate emissions associated with the electricity supply sector to each of the RCI sectors. Electricity sales associated with the RCI sectors were estimated based on information from the *Patterns and Trends* report¹³

1.3 Results

1.3.1 Residential Sector

The emissions inventory for the Residential sector presented in Figure 1-1 was developed from the emissions data in Table 1-2. Table 1-3 shows the relative contributions of emissions associated with each fuel type to total Residential sector emissions. For the Residential sector, emissions from direct fuel use (including heating and hot water) and electricity consumption decrease by 11 percent, from approximately 54 MMtCO₂e in 1990 to 48 MMtCO₂e in 2014. Emissions associated with natural gas consumption were the greatest source of emissions in 2014, accounting for approximately 52 percent. Emissions associated with electricity use were the second largest source, with 27 percent of the total in this sector. Petroleum use accounted for the next largest portion of Residential sector emissions at 21 percent of emissions. Coal and wood emissions account for one percent or less of Residential sector emissions throughout the analysis period. The Residential sector's share of total RCI emissions from direct fuel use and electricity increased from 37 percent in 1990 to 46 percent in 2014.

¹³ NYSERDA. 2016. "Patterns and Trends, New York State Energy Profiles: 2000-2014," (<https://www.nyserda.ny.gov/About/Publications/EA-Reports-and-Studies/Patterns-and-Trends>)

Figure 1-1. Residential Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014

The Other category includes emissions associated with coal and wood combustion. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted (i.e., the CO₂ emissions are zero), as per EPA guidance. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.




Table 1-2. Residential Sector Emissions Inventory, 1990–2014 (MMtCO₂e)

The Other category includes emissions associated with coal and wood combustion. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted (i.e., the CO₂ emissions are zero), as per EPA guidance. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.

Fuel Type	1990	1995	2000	2005	2010	2014
Other	0.5	0.5	0.7	0.4	0.1	0.2
Petroleum	15.3	13.9	17.6	17.2	10.3	10.2
Natural Gas	18.5	20.6	22.0	22.2	21.2	25.1
Electricity	19.3	16.4	18.6	20.2	16.5	13.0
Total	53.5	51.4	58.8	60.0	48.2	48.5

Table 1-3. Residential Sector Proportions of Total Emissions by Fuel Type, 1990–2014 (%)


The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table 1-2

Fuel Type	1990	1995	2000	2005	2010	2014
Other	0.9%	1.0%	1.2%	0.7%	0.3%	0.4%
Petroleum	28.6%	27.0%	29.9%	28.7%	21.4%	21.0%
Natural Gas	34.6%	40.0%	37.3%	37.0%	44.1%	51.8%
Electricity	36.0%	32.0%	31.6%	33.6%	34.3%	26.8%

The trend in GHG emissions from fuel combustion for the Residential sector, excluding emissions associated with electricity use, correlates with the trend in the number of heating degree days for each year from 1990–2014, as demonstrated in the comparison in Figure 1-2.¹⁴ The two sets of data show a correlation coefficient of 0.60. Although the magnitude of the effect of heating degree days on emissions varies over time, the important point to note is that the peaks and valleys of both sets of data occur in the same years. In years with a greater number of heating degree days, the emissions are also generally higher, and in years with a low number of heating degree days, such as 1998 and 2006, GHG emissions from Residential sector were also lower.

¹⁴ Heating degree days provide a measurement of how cold a location is relative to a base temperature, which is typically 65° F, over a period of time. The measure is calculated by subtracting the average of a day's high and low temperatures from the base temperature, with negative values set to equal zero.

Figure 1-2. Residential Sector Heating Demand and GHG Emissions, 1990–2014



1.3.2 Commercial/Institutional Sector

The emissions inventory for the Commercial/Institutional sector presented in Figure 1-3 was developed from the emissions data in Table 1-4. Table 1-5 shows the relative contributions of emissions associated with each fuel type to total Commercial/Institutional sector emissions.

For the Commercial/Institutional sector, emissions from direct fuel use (including heating and hot water) and electricity decreased from approximately 54 MMtCO₂e in 1990 to 42 MMtCO₂e in 2014. Electricity use was the largest source of all Commercial/Institutional sector emissions at 51 percent in 1990, decreasing slightly to 48 percent by 2014. Petroleum emissions accounted for 28 percent of Commercial/Institutional sector emissions in 1990, declining to 11 percent in 2014. In 1990, natural gas consumption accounted for approximately 20 percent of total commercial/institutional emissions and increased to 42 percent by 2014. Commercial/Institutional sector emissions associated with the use of coal accounted for approximately one percent of total commercial/institutional emissions in 1990 and less than one percent of total commercial/institutional emissions in 2014. The Commercial/Institutional sector's share of total RCI emissions from direct fuel use and electricity use was 38 percent in 1990 and 39 percent in 2014.

Figure 1-3. Commercial/Institutional Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014

The Other category includes emissions associated with coal and wood combustion. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted (i.e., the CO₂ emissions are zero), as per EPA guidance. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.




Table 1-4. Commercial/Institutional Sector Emissions Inventory, 1990–2014 (MMtCO₂e)

The Other category includes emissions associated with coal and wood combustion. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted (i.e., the CO₂ emissions are zero), as per EPA guidance. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.

Fuel Type	1990	1995	2000	2005	2010	2014
Other	0.5	0.5	0.2	0.4	0.0	0.0
Petroleum	15.2	13.8	11.8	13.1	8.5	4.4
Natural Gas	10.6	12.7	20.0	15.0	15.6	17.5
Electricity	28.0	25.8	30.4	30.6	25.0	20.0
Total	54.4	52.7	62.5	59.1	49.1	41.9

Table 1-5. Commercial/Institutional Sector Proportions of Total Emissions by Fuel Type (%)

The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table 1-4.

Fuel Type	1990	1995	2000	2005	2010	2014
Other	1.0%	0.9%	0.4%	0.6%	0.0%	0.0%
Petroleum	28.0%	26.1%	18.8%	22.2%	17.3%	10.6%
Natural Gas	19.6%	24.0%	32.1%	25.4%	31.7%	41.8%
Electricity	51.4%	48.9%	48.7%	51.8%	51.0%	47.6%

1.3.3 Industrial Sector

The emissions inventory for the Industrial sector presented in Figure 1-4 was developed from the emissions data in Table 1-6. Table 1-7 shows the relative contributions of emissions associated with each fuel type to total Industrial sector emissions.

For the Industrial sector, emissions from direct fuel use and electricity in 1990 were approximately 36 MMtCO₂e, but decreased to approximately 16 MMtCO₂ in 2014. In 1990, the emissions associated with electricity use accounted for the largest share of industrial emissions, at 44 percent of the Industrial sector's total. However, this contribution declines to 30 percent in 2014. In contrast, emissions from petroleum fuels accounted for 18 percent of the Industrial sector emissions in 1990, but account for 20 percent of industrial emissions in 2014. The share of emissions from natural gas consumption increased from 16 percent of industrial fuel use emissions in 1990 to 38 percent of these emissions in 2014. Coal consumption accounted for approximately 22 percent of total industrial emissions in 1990 and decreased to approximately 13 percent of total industrial emissions in 2014. The Industrial sector's share of total RCI emissions from direct fuel use and electricity use was 25 percent in 1990 and decreased to 15 percent in 2014.

Figure 1-4. Industrial Sector GHG Emissions from Fuel Combustion and Electricity, 1990–2014

Emissions associated with wood combustion are too small to be seen on this graph. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted (i.e., the CO₂ emissions are zero), as per EPA guidance. Therefore, the CO₂ emissions are zero. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.




Table 1-6. Industrial Sector Emissions Inventory, 1990–2014 (MMtCO₂e)

The Other category includes emissions associated with wood combustion. Wood is considered a biomass fuel, and carbon dioxide emissions from biomass fuels grown sustainably are not counted. Therefore, the CO₂ emissions are zero, as per EPA guidance. Consequently, the GHG emissions associated with this category include methane and nitrous oxide. In general, GHG emissions include all six standard GHGs, expressed in MMtCO₂e.

Fuel Type	1990	1995	2000	2005	2010	2014
Coal	7.7	6.3	6.5	3.7	2.3	2.0
Petroleum	6.5	4.2	5.3	6.4	3.1	3.1
Natural Gas	5.7	11.9	5.6	4.9	4.8	5.9
Electricity	16.0	12.1	11.2	8.2	4.3	4.7
Total	35.9	34.6	28.6	23.2	14.6	15.7

Table 1-7. Industrial Sector Proportions of Total Emissions by Fuel Type (%)

The percentages shown in this table reflect the emissions for each fuel type as a percentage of total emissions shown in Table 1-5.

Fuel Type	1990	1995	2000	2005	2010	2014
Coal	21.6%	18.3%	22.8%	15.7%	16.0%	12.7%
Petroleum	18.0%	12.1%	18.5%	27.6%	21.0%	20.0%
Natural Gas	15.9%	34.4%	19.5%	21.1%	33.1%	37.5%
Electricity	44.4%	35.1%	39.0%	35.5%	29.8%	29.8%

2 Industrial Non-Fuel Combustion Processes

2.1 Overview

Emissions from industrial non-fuel combustion sources span a wide range of industrial processes and energy supply activities. The following sections describe those processes and activities that exist in the State and their estimated emissions in this inventory.

2.1.1 Sources of Carbon Dioxide (CO₂) Emissions

Cement production: Greenhouse gas (GHG) emissions related to cement production can come from both clinker and cement kiln dust. Clinker is an intermediate product used to make finished Portland and masonry cement. Clinker production releases CO₂ when calcium carbonate is heated in a cement kiln to form lime (calcium oxide) and CO₂.¹⁵ About 0.02 metric tons (Mt) of CO₂ is emitted for every Mt of cement kiln dust produced, relative to the CO₂ emitted during the production of an Mt of clinker.¹⁶

Iron and steel production: The production of iron and steel generates process-related CO₂ emissions. The process of reducing iron ore with metallurgical coke in a blast furnace to create pig iron, which is then used as a raw material in the production of steel, emits CO₂. The production of metallurgical coke from coking coal produces CO₂ emissions as well.

Limestone and dolomite use: Limestone and dolomite are basic raw materials used by a wide variety of industries, including the construction, agriculture, chemical, glass manufacturing, and environmental pollution control industries as well as metallurgical industries such as magnesium production. The industrial processes sector also includes the emissions associated with the use of limestone and dolomite to manufacture steel and glass, and their use in flue-gas desulfurization scrubbers to control sulfur dioxide emissions from the combustion of coal in boilers.¹⁷

¹⁵ For further detail, see EIIP. 2006. "Chapter 6: Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII.

¹⁶ EIIP. 2006. "Chapter 6: Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII.

¹⁷ In accordance with EIIP Chapter 6 methods, emissions associated with the following uses of limestone and dolomite are not included in this category: (1) crushed limestone consumed for road construction or similar uses (because these uses do not result in CO₂ emissions), (2) limestone used for agricultural purposes (which is counted under the methods for the agricultural sector), and (3) limestone used in cement production (which is counted in the methods for cement production).

Soda ash use: Commercial soda ash (sodium carbonate) is found in many consumer products such as glass, soap and detergents, paper, textiles, and food. Consuming soda ash releases carbon dioxide.¹⁸

Other industrial processes that produce CO₂ emissions, but are not found in New York State or for which reliable or complete data are lacking, are taconite and lime production and ammonia production. In addition, CO₂ emissions can also result from urea consumption. While the SIT provides default data for urea consumption in the State, the estimated amount of emissions from this source is small and is excluded from the inventory.¹⁹

2.1.2 Sources of Methane (CH₄) Emissions

Natural gas leakage: Methane emissions are associated with the transmission, storage, and distribution of natural gas in the State.²⁰ Transmission pipelines are large diameter, high-pressure lines that transport gas from production fields, processing plants, storage facilities, and other sources of supply over long distances to local distribution companies or to large volume customers. Sources of CH₄ emissions from transmission pipelines include leaks, compressor fugitives, vents, and pneumatic devices. Distribution pipelines are extensive networks of generally small diameter, low-pressure pipelines that distribute gas within cities or towns. Sources of CH₄ emissions from distribution pipelines include leaks, meters, and regulators.

¹⁸ For further detail, see EIIP. 2006. "Chapter 6 Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII.

¹⁹ The default SIT data indicate that CO₂ emissions from urea consumption amounted to <0.005 MMtCO₂e annually between 1990 and 2014.

²⁰ Note that CH₄, N₂O, and CO₂ emissions from natural gas consumed as lease fuel (used in well, field, and lease operations), plant fuel (used in natural gas processing plants), and pipeline fuel (used in pipeline compressor station internal combustion engines) are included in Chapter 1 in the industrial fuel combustion category.

New York State imports the vast majority of its natural gas via pipeline from other states and Canada. The Transcontinental and Tennessee Gas Transmission pipelines from the Gulf Coast and the Iroquois pipeline from Canada link up with local gas distribution networks that supply the New York City metropolitan area and Long Island. Other parts of the State receive gas from Pennsylvania and Canada via other gas transmission systems.

2.1.3 Sources of Perfluorocarbon (PFC) Emissions

Aluminum production: Emissions of tetrafluoromethane and hexafluoroethane, both PFCs, occur during the reduction of alumina in the primary smelting process. Based on EPA's SIT methodology, the aluminum production industry is the largest source of these two PFCs.

*Semiconductor manufacturing:*²¹ Manufacturers of semiconductors use fluorinated GHGs in the plasma etching and plasma enhanced chemical vapor deposition processes. Plasma etching of dielectric films creates the pattern of pathways connecting individual circuit components in semiconductors. Vapor deposition chambers used for depositing the dielectric films are periodically cleaned using fluorinated gases. Converting fluorinated gases to fluorine atoms in plasma etches away dielectric material or cleans the chamber walls and hardware. Un-dissociated fluorinated gases and other products end up in the waste streams and, unless captured by abatement systems, into the atmosphere. Some fluorinated compounds can transform into other compounds (e.g., CF₄ generated from C₂F₆) during the plasma processes. If these other compounds are not captured by emission control systems, the process-generated gases are also released into the atmosphere.²²

²¹ Emissions from semiconductor manufacturing were estimated using the EPA's State Greenhouse Gas Inventory Tool (SIT) software, with reference to EIIP. 2006. "Chapter 6 Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII. Though SIT provides an aggregate emissions value for PFCs, hydrofluorocarbons, and sulfur hexafluoride, it is assumed PFCs constitute the majority of the emissions.

²² California Environmental Protection Agency, Air Resources Board. 2009. California's 1990–2004 Greenhouse Gas Emissions Inventory and 1990 Emissions Level: Technical Support Document.

2.1.4 Sources of Hydrofluorocarbon (HFC) Emissions

Ozone-depleting substances (ODS) substitutes: HFC emissions result from the consumption of substitutes for ozone-depleting substances used in cooling and refrigeration equipment.²³ The most notable HFC substitution is for CFCs, which are also potent warming gases that have global warming potentials on the order of thousands of times that of CO₂ per unit of mass. This substitution is in compliance with the Montreal Protocol and the Clean Air Act. Amendments of 1990.²⁴ In October of 2016, the Montreal Protocol was amended to limit HFC usage. Even low amounts of HFC emissions (e.g., from leaks and other releases associated with normal use of the products) can lead to high GHG emissions on a CO₂e basis. HCFC-22 production is another industrial process that can result in HFC emissions. However, specific data for New York State in this source category are lacking.

2.1.5 Other Sources of Non-Fuel Combustion Emissions

Nitrous Oxide Emissions: Nitric acid production is an industrial process that can result in nitrous oxide emissions. However, specific data for New York State in this source category are lacking.

Sulfur Hexafluoride Emissions: Magnesium production and processing are industrial processes that can result in sulfur hexafluoride emissions. However, specific data for New York State in this source category are lacking.

2.2 Emissions Inventory Data and Methodology

GHG emissions for 1990 through 2014 were estimated using the following methods. For most sources, emissions were estimated using the SIT tool and the methods provided in the EIIP guidance document for industrial processes.²⁵

²³ Environmental Protection Agency. 2016. Draft User's Guide for Estimating Carbon Dioxide, Nitrous Oxide, HFC, PFC, and SF₆ Emissions from Industrial Processes Using the State Industrial Tool. Substitutes for ozone-depleting substances, which include chlorofluorocarbons, halons, carbon tetrachloride, methyl chloroform, and hydrochlorofluorocarbons, are used in a variety of industrial applications including refrigeration and air conditioning equipment, aerosols, solvent cleaning, fire extinguishing, foam blowing, and sterilization. Although their substitutes, HFCs, are not harmful to the stratospheric ozone layer, they are powerful GHGs.

²⁴ As noted in EIIP Chapter 6, ODS substitutes are primarily associated with refrigeration and air conditioning, but also many other uses including as fire control agents, cleaning solvents, aerosols, foam blowing agents, and in sterilization applications. The applications, stocks, and emissions of ODS substitutes depend on technology characteristics in a range of equipment types. For the U.S. national inventory, a detailed stock vintaging model was used to track ODS substitutes uses and emissions, but this modeling approach has not been completed at the state level.

²⁵ GHG emissions were calculated using SIT, with reference to EIIP. 2006. "Chapter 6: Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII.

2.2.1 Sources of CO₂ Emissions

Cement production: Emissions from cement production are calculated using the SIT methodology by multiplying annual metric tons of clinker production by emission factors to estimate emissions associated with the clinker production process (0.507 Mt of CO₂ emitted per Mt of clinker produced) and cement kiln dust (0.020 MtCO₂ emitted per Mt of clinker CO₂ emitted). Data on the Mt of clinker produced in the State were derived from the cement statistics in United States Geological Survey's (USGS) Minerals Yearbook.²⁶ This information is available within the SIT module.

Iron and steel production: New York State's emissions from iron and steel production were pro-rated from the EPA's estimates of total U.S. emissions²⁷ based on the State's market share of national iron and steel manufacturing derived from American Iron and Steel (AISI) data.²⁸ The basic activity data used are the quantities of crude steel produced (defined as first cast product suitable for sale or further processing). This information is available within the SIT module as default. For inventory years 2011–2014, an annual average of data from production facilities reported to EPA through its *Facility Level Information on GreenHouse gases* Tool (FLIGHT) was used.

Limestone use: Historical State emissions were pro-rated from the EPA's estimates of total U.S. emissions from limestone and dolomite consumption²⁹ based on the quantity of crushed stone sold in the State derived from USGS data.³⁰ This information is available within the SIT module as default.

²⁶ U.S. Geological Survey. 2016. "Minerals Yearbook: Cement Statistics and Information" Last modified 15 November 2016. <http://minerals.usgs.gov/minerals/pubs/commodity/cement/index.html#myb>. The metric tons of cement clinker produced in New York State are assumed to be half of the total production value for Maine and New York.

²⁷ Environmental Protection Agency. 2016. "Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2014." <https://www.epa.gov/ghgemissions/us-greenhouse-gas-inventory-report-1990-2014>.

²⁸ NY's Market Share of U.S. Iron & Steel Manufacture was prorated from national data provided by the American Iron and Steel (AISI) Annual Statistics Report 2009 (AISI 2011) within the SIT Module.

²⁹ Environmental Protection Agency. 2016. "Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2014." <https://www.epa.gov/ghgemissions/us-greenhouse-gas-inventory-report-1990-2014>.

³⁰ U.S. Geological Survey. 2016 "Minerals Yearbook Crushed Stone Statistics and Information." Last modified 17 August 2016. http://minerals.usgs.gov/minerals/pubs/commodity/stone_crushed/.

Soda ash use: Emissions from soda ash use are calculated within the SIT module by scaling national soda ash consumption (estimated using sales data) based on the ratio of state to national population.³¹ Data on the metric tons of soda ash consumed in the U.S. were derived from the soda ash statistics in the USGS Minerals Yearbook.³² This information is available within the SIT module as default.

2.2.2 Sources of CH₄ Emissions

Natural gas leakage: The EPA published estimates³³ of national CO₂e emissions for natural gas systems for 1990 through 2014.³⁴ These estimates are reported for four processes:

- Production.
- Processing.
- Transmission and storage.
- Distribution.

Natural gas leakage in the State was analyzed for the last two processes, given the limited occurrence of production and processing in New York State. Emissions from natural gas transmission, storage, and distribution were estimated by scaling the U.S. GHG emissions from these processes based on the ratio of the State's natural gas consumption to national natural gas consumption. The State's natural gas consumption estimates for 2014 were obtained from NYSERDA's *Patterns and Trends* report³⁵ and national consumption estimates were compiled from Energy Information Administration (EIA) State Energy Data System (SEDS) data.³⁶ The science underlying methane emissions calculation is advancing quickly, and methods of estimating emissions may require reevaluation and refinement going forward in accordance with evolving standards and best practices.

³¹ U.S. Census Bureau. 2010. <http://www.census.gov/2010census/>

³² U.F S. Geological Survey. 2016. "Minerals Yearbook: Soda Ash Statistics and Information." Last modified 15 November 2016. http://minerals.usgs.gov/minerals/pubs/commodity/soda_ash/.

³³ A recent study found that measured emissions of methane from natural gas systems are approximately 1.5 times greater than those published in the U.S. Environmental Protection Agency national GHG inventory. A commensurate scaling of this analysis would increase the emissions from natural gas leakage to 3.3 MMTonsCO₂e or 1% of total emissions. See Brandt, et al. 2014. "Methane Leaks from North American Natural Gas Systems." *Science* 343, February.

³⁴ Environmental Protection Agency. 2016. "Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014." (<https://www.epa.gov/ghgemissions/us-greenhouse-gas-inventory-report-1990-2014>)

³⁵ NYSERDA. 2016. "Patterns and Trends, New York State Energy Profiles: 1999-2014." <https://www.nyserdera.ny.gov/About/Publications/EA-Reports-and-Studies/Patterns-and-Trends>.

³⁶ U.S. Energy Information Administration. 2016. "State Energy Data System 1960-2014." <http://www.eia.gov/state/seds/seds-data-complete.cfm?sid=US>.

2.2.3 Sources of PFC Emissions

Aluminum production: Emissions from aluminum production are calculated using the SIT Industrial Processes module methodology based on the annual quantity of aluminum production. Data on the metric tons of aluminum produced in New York were derived from the aluminum statistics in the USGS Minerals Yearbook.³⁷

Semiconductor manufacturing: Historical emissions associated with semiconductor manufacturing were estimated using the SIT calculation methodology. The SIT calculates these emissions based on Economic Census data indicating the State's portion of the national dollar value of semiconductor shipments along with estimates of national emissions per shipment dollar from the EPA national GHG inventory.

2.2.4 Sources of HFC Emissions

ODS substitutes: Historical emissions from ODS substitutes are calculated within the SIT by scaling national emissions from the EPA's GHG inventory³⁸ based on the ratio of state to national population.³⁹

2.3 Results

Figure 2-1 shows emissions from industrial non-fuel combustion processes from 1990–2014, which are based on the historical emission values in Table 2-1. Total State GHG emissions for these sources totaled approximately 8.2 MMtCO₂e in 1990 and grew to 14.1 MMtCO₂e in 2014. Emissions growth is primarily associated with the increasing use of ozone-depleting substances (ODS) substitutes, as shown in Figure 2-1.

³⁷ U.S. Geological Survey. 2016. "Minerals Yearbook, Aluminum Statistics and Information." Last modified 25 October 2016. <http://minerals.usgs.gov/minerals/pubs/commodity/aluminum/>.

³⁸ Environmental Protection Agency. 2016. "Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990–2014." <https://www.epa.gov/ghgemissions/us-greenhouse-gas-inventory-report-1990-2014>.

³⁹ U.S. Census Bureau. 2010. <http://www.census.gov/2010census/>

Figure 2-1. GHG Emissions from Industrial Non-Fuel Combustion Processes by Source, 1990–2014




Table 2-1. Emissions Inventory for Industrial Non-Fuel Combustion Processes, 1990–2014 (MMtCO₂e)

Industrial Process	1990	1995	2000	2005	2010	2014
CO₂ Emissions						
Cement Manufacture	0.7	0.8	0.8	0.8	0.5	0.4
Iron & Steel Production	0.8	0.8	0.8	0.6	0.2	0.1
Limestone & Dolomite Use	0.2	0.3	0.2	0.4	0.4	0.6
Soda Ash	0.2	0.2	0.2	0.2	0.1	0.1
CH₄ Emissions						
Natural Gas Systems	4.6	5.2	3.9	2.6	2.0	2.2
PFC Emissions						
Aluminum Production	0.4	0.3	0.3	0.3	0.3	0.2
Semiconductor Manufacturing	0.1	0.1	0.1	0.1	0.1	0.1
HFC Emissions						
ODS Substitutes	0.0	2.4	6.0	7.3	9.1	10.0
Electric Power Transmission and Distribution Systems	1.2	0.9	0.6	0.4	0.3	0.2
TOTAL	8.2	11.0	12.8	12.6	13.1	14.1

2.3.1 Sources of CO₂ Emissions

Cement manufacturing: Emissions estimates from this source were approximately 0.7 MMtCO₂e in 1990, decreasing slightly to 0.4 MMtCO₂e in 2014, as shown in Figure 2-1 and Table 2-1.

Iron and steel production: Emissions in 1990 were 0.8 MMtCO₂e and decreased to about 0.1 MMtCO₂e in 2014, as shown in Figure 2-1 and Table 2-1.

Limestone and Dolomite use: Emissions from limestone consumption were approximately 0.2 MMtCO₂e in 1990 and 0.6 MMtCO₂e in 2014.

Soda ash use: Emissions estimates from soda ash consumption were approximately 0.2 MMtCO₂e in 1990 and 0.1 MMtCO₂e in 2014.

2.3.2 Sources of CH₄ Emissions

Natural gas leakage: Natural gas leakage is currently the second highest contributor to GHG emissions from industrial non-fuel combustion processes in the State. Emissions estimates from this source were 4.6 MMtCO₂e in 1990 and 2.2 MtCO₂e in 2014.⁴⁰ This methane emissions trajectory is methodologically tied to EPA's emissions assumptions from natural gas transmission, distribution, and storage, and driven by EPA's assumptions that emissions from these elements of the natural gas system have declined from 1990 through 2014. Note that EPA has revised its national estimates of fugitive methane emissions from natural gas extraction sites from 1990–2014 higher to better reflect current scientific research on methane leakage. These emissions are assumed to occur outside of the State. The science underlying methane emissions calculation is advancing quickly, and methods of estimating emissions may require reevaluation and refinement going forward in accordance with evolving standards and best practices.

2.3.3 Sources of PFC Emissions

Semiconductor manufacturing: Emissions from this source are fairly low, at 0.1 MMtCO₂e, in New York throughout the analytic period, as shown in Figure 2-1 and Table 2-1.

Aluminum production: GHG emissions estimates from aluminum production in the State were 0.4MMtCO₂e and 0.2 MMtCO₂e in 2014.

2.3.4 Sources of HFC Emissions

ODS substitutes: Estimates of emissions from ODS substitutes in New York State have increased from 0.02 MMtCO₂e in 1990 to about 10 MMtCO₂e in 2014. ODS substitutes (and as a result, HFCs) have increased due to the provisions of the Montreal Protocol. While the protocol was effective at protecting the ozone layer by phasing out ODSs, it has led to an increase in HFCs, which are a potent GHG. Given the risk HFCs pose to the climate, the EPA has enacted new rules to limit emissions from HFCs under its Significant New Alternatives Policy.⁴¹ More recently, the Montreal Protocol was amended in October 2016 to limit HFC usage.

⁴⁰ A recent study found that measured emissions of methane from natural gas systems are approximately 1.5 times greater than those published in the U.S. Environmental Protection Agency national GHG inventory. A commensurate scaling of this analysis would increase the emissions from natural gas leakage to 3.3 MMTonsCO₂e or 1% of total emissions. See Brandt, et al. 2014. "Methane Leaks from North American Natural Gas Systems." *Science* 343, February.

⁴¹ Environmental Protection Agency. 2016. "Significant New Alternatives Policy." Last modified 11 October 2016. <https://www.epa.gov/snap/snap-regulations>.

3 Transportation Energy Use

3.1 Overview

The transportation sector is one of the largest sources of GHG emissions in the State. Carbon dioxide (CO₂) accounts for the majority of the transportation GHG emissions from fuel use. Most of the remaining GHG emissions from the transportation sector are due to nitrous oxide (N₂O) emissions from on-road gasoline-fueled vehicles.

3.1.1 On-Road Vehicle Fuels

Gasoline: Gasoline combusted by passenger cars, light-duty trucks, heavy-duty vehicles, and motorcycles produces CO₂, N₂O, and CH₄ emissions. Although gasoline contains virtually no CH₄, this GHG is a byproduct of gasoline combustion and is influenced by various factors such as emission control technologies, combustion conditions, and fuel composition. Gasoline contains up to 10 percent ethanol by volume; however, per the EPA's EIIP guidance documents⁴² ethanol is considered a biofuel, so no CO₂ emissions associated with the combustion of ethanol blended into gasoline are included in the inventory.⁴³

Diesel: Distillate fuel combusted by passenger cars, light-duty trucks, and heavy-duty vehicles produces CO₂, N₂O, and CH₄ emissions. Although diesel contains virtually no CH₄, it is a byproduct of diesel combustion and is influenced by various factors such as emission control technologies, combustion conditions, and fuel composition. To the extent that diesel fuel is blended with biodiesel, CO₂ emissions from this biofuel are not included in the inventory per EPA guidance.

Natural gas: Natural gas combusted by light-duty and heavy-duty alternative fuel vehicles produces CO₂, N₂O and CH₄ emissions.⁴⁴

⁴² EIIP. 2006. "Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels." Volume VIII.

⁴³ Carbon dioxide emissions associated with fuels produced from sustainably grown biomass are not counted. For cases where biomass is not grown sustainably, the GHG impact would be captured as a land use change.

⁴⁴ Note that natural gas use for transportation application is often called 'compressed natural gas' or 'CNG' in other state and federal documents and databases.

3.1.2 Off-Road Vehicle Fuels

Gasoline: CO₂, N₂O, and CH₄ emissions are produced from gasoline combusted by sources such as marine vehicles.

Diesel: CO₂, N₂O, and CH₄ emissions are produced from distillate fuel combusted by such sources as locomotives and marine vehicles.

Residual oil: CO₂, N₂O, and CH₄ emissions are produced from residual fuel combusted by marine vehicles.

Liquefied petroleum gas: CO₂, N₂O, and CH₄ emissions are produced from LPG combusted by internal combustion engines.⁴⁵

Jet fuel (kerosene): CO₂, N₂O, and CH₄ emissions are produced from jet fuel combusted by aircraft.

3.1.3 Lubricants

CO₂ emissions are produced from lubricants that are consumed during vehicle operation. Lubricants have an assumed oxidation factor of 1.00,⁴⁶ which represent the fraction of consumed product (on an energy-content basis) that is combusted and leads to GHG emissions.

⁴⁵ U.S. Energy Information Administration. State Energy Data System Technical Notes and Documentation, Consumption, Section 4: Petroleum. http://www.eia.gov/state/seds/sep_use/notes/use_petrol.pdf.

The LPG data from the United States Department of Energy's Energy Information Administration (EIA) captures LPG sales for the internal combustion engines of highway vehicles, forklift, industrial tractors, and for use in oil field drilling and production. It is assumed that New York State has very few highway vehicles that consume LPG, therefore all LPG data is attributed to off-road vehicles.

⁴⁶ The oxidation factor for lubricants was derived from EPA's State Inventory Tool CO2FFC Module, with reference to EIIP, 2006. "Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels." Volume VIII.

3.2 Emissions Inventory Data and Methodology

Historical GHG emissions estimates use the SIT and the methods provided in the EIIP guidance document for the sector.^{47,48}

The SIT module was used to determine emissions from the following sources: Highway vehicles, aviation, boat and marine vessels, alternative fuel vehicles. Options for locomotives and “other non-highway sources” were not utilized. Default emissions factors were used.

3.2.1 On-Road Vehicle Fuels

The SIT CO₂ Fossil Fuel Combustion Module was used to calculate CO₂ emissions based on the energy content, in British thermal units (Btu), of fuel and lubricants consumed. The default SIT fuel consumption values for the State were replaced with State-specific information calculated from annual vehicle miles of travel (VMT) data provided by NYS Department of Transportation (DOT) and fuel economy data provided by Oak Ridge National Laboratory (ORNL) and Federal Highways Administration (FHWA). The SIT Mobile Combustion Module was used to calculate N₂O and CH₄ emissions based VMT for on-road vehicles, with the SIT default data for VMT replaced with DOT data for New York State.

Gasoline and diesel: New York State annual VMT data for gasoline and diesel on-road vehicles were allocated to differing vehicle types to calculate the VMT by vehicle type for 1990, 2002, 2007, and 2014 shown in Table 3-1.⁴⁹ The fraction of VMT by vehicle type for years 1991–2001, 2002–2007, and 2008–2014 were calculated by interpolation and the fractions for years after 2014 were held constant at the 2011 levels. These fractions, applied to the corresponding annual statewide total VMT value to produce annual VMT by vehicle type, were input into the SIT Mobile Combustion Module to calculate emissions of N₂O and CH₄.

⁴⁷ CO₂ emissions were calculated using SIT, with reference to EIIP. 2006. “Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels.” Volume VIII.

⁴⁸ CH₄ and N₂O emissions were calculated using SIT, with reference to EIIP. 2006. “Chapter 3 Methods for Estimating Methane and Nitrous Oxide Emissions from Mobile Combustion.” Volume VIII.

⁴⁹ New York State Department of Transportation VMT modeling.

Table 3-1. New York State VMT by Vehicle Type (million miles)

Vehicle Type	1990	2002	2007	2014
Heavy Duty Diesel Vehicle	4,056	4,538	8,022	4,255
Heavy Duty Gasoline Vehicle	206	320	1,414	1,120
Light Duty Diesel Truck	582	1,890	1,459	1,119
Light Duty Diesel Vehicle	26	20	19	18
Light Duty Gasoline Truck	17,552	61,268	64,119	60,321
Light Duty Gasoline Vehicle	84,372	64,420	60,817	59,987
Motorcycle	107	603	887	4,080
Total	106,900	133,058	136,737	130,889

Annual fuel efficiency data were applied to the annual VMT by vehicle type to calculate the quantity of on-road fuel consumption. ORNL and FHWA provided the historical fuel efficiency data. Consumption of biofuel within blended fuels was estimated by applying ethanol fractions of motor gasoline and biodiesel fractions of diesel sold to the calculated consumption numbers. The biofuel fractions were obtained from EIA's State Energy Data System (SEDS) for New York State. The estimated consumption of the biofuels was then subtracted from the gasoline and diesel totals and the resultant consumption data were input into the SIT CO₂FFC Module to calculate emissions of CO₂.

Natural Gas: Historical transportation natural gas consumption data obtained from EIA's SEDS were input into the CO₂FFC Module to calculate CO₂ emissions.

Consumption data from EIA's SEDS and fuel efficiency data from ORNL and FHWA provided the base for the State's annual natural gas historical VMT data by vehicle type calculations. The resultant VMT values were then input into the SIT Mobile Combustion Module to calculate emissions of N₂O and CH₄.

3.2.2 Off-Road Vehicle Fuels

Gasoline: Estimates for historical off-road motor gasoline consumption are based on FHWA marine vehicle gasoline sales data for New York State. These data were divided by the FHWA national motor gasoline sales data to calculate the fraction of sales associated with New York State marine vehicles. These fractions scale the national motor gasoline consumption data from SEDS to estimate New York State off-road motor gasoline consumption. Gasoline consumption data in Btu were subsequently input into the SIT CO₂FFC and Mobile Combustion Modules to produce GHG emissions estimates.

Diesel: For historical analysis, estimates for off-road diesel consumption use the off-road diesel sales data from SEDS. The diesel sales data were then adjusted using national data on the ratio of diesel consumption to diesel sales in the U.S. in order to approximate off-road diesel consumption. Diesel consumption data in Btu were subsequently input into the SIT CO₂FFC and Mobile Combustion Modules to produce GHG emissions estimates.

Residual oil, liquefied petroleum gas, and jet fuel (kerosene): For these other off-road fuels, emissions estimates were calculated within the SIT CO₂FFC and Mobile Combustion modules using fuel consumption data in Btu from NYSERDA's *Patterns and Trends* report.⁵⁰

3.2.3 Lubricants

Historical emissions estimates were calculated using fuel consumption data in Btu from NYSERDA's *Patterns and Trends* and the historical CO₂ emission estimates were calculated within the SIT CO₂FFC module.

3.3 Results

Gasoline consumption accounts for the largest share of transportation GHG emissions, as shown in Table 3-2 and Figure 3-1. From 1990–2014, emissions from gasoline increased by about 4 percent to account for approximately 73 percent of total transportation emissions in 2014. GHG emissions from diesel fuel remained consistent from 1990–2014, accounting for approximately nine percent of total transportation emissions in 2014. Emissions from jet fuel also accounted for approximately 11 percent of transportation emissions in 2014. Emissions from residual oil accounted for about five percent of total transportation emissions in 2014, while emissions from all other categories combined (natural gas, LPG, and lubricants) contributed approximately one percent of total transportation emissions in 2014.

⁵⁰ NYSERDA. 2016. "Patterns and Trends, New York State Energy Profiles: 2000–2014," (<https://www.nyserdera.ny.gov/About/Publications/EA-Reports-and-Studies/Patterns-and-Trends>)

Figure 3-1. Transportation GHG Emissions by Fuel, 1990–2014

LPG is liquefied petroleum gas.




Table 3-2. Transportation Sector Emissions Inventory, 1990–2014 (MMtCO₂e)

Fuel Type	1990	1995	2000	2005	2010	2014
Gasoline	50.62	53.43	59.44	61.97	54.38	54.25
Jet Fuel, Kerosene	1.54	2.19	2.75	5.81	4.24	8.32
Diesel	7.07	7.36	9.03	12.66	9.96	7.06
Residual Oil	0.65	1.09	3.87	2.70	5.68	3.60
Lubricants	0.48	0.46	0.49	0.42	0.39	0.36
LPG	0.04	0.03	0.06	0.02	0.03	0.08
Natural Gas	0.00	0.02	0.05	0.15	0.23	0.34
Total	60.40	64.57	75.68	83.73	74.91	74.01

4 Power Supply and Delivery

4.1 Overview

The generation, transmission, and distribution of electric power to meet electricity demand in New York State produces carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), and sulfur hexafluoride (SF₆) emissions. The power supply and delivery sector GHG emission sources and sinks are subdivided into three primary categories: fuel combustion, electricity distribution, and municipal waste combustion.

4.1.1 Fuel Combustion

Electricity generation: This source category captures CO₂, CH₄, and N₂O emissions that result from the burning of various fuels by in-State power plants.

Net imports of electricity: New York State has historically imported more electricity than it has exported. Net imports are the difference between total in-State imports and exports. This source category includes CO₂, CH₄, and N₂O emissions associated with net imported electricity.

4.1.2 Electricity Distribution

This source category captures SF₆ emissions associated with the transmission and distribution (T&D) of electricity. SF₆ is used as an electrical insulator and interrupter in the electric power T&D system. Because of its high dielectric strength and arc-quenching abilities, the largest use for SF₆ is as an insulator in T&D equipment such as gas-insulated high-voltage circuit breakers, substations, transformers, and transmission lines. Not all of the electric utilities in the U.S. use SF₆; use of this gas is more common in urban areas where the space occupied by electric power T&D facilities is more valuable.⁵¹

⁵¹ Environmental Protection Agency. 2013 User's Guide for Estimating Carbon Dioxide, Nitrous Oxide, HFC, PFC, and SF₆ Emissions from Industrial Processes Using the State Inventory Tool. Prepared by ICF International.

4.1.3 Municipal Waste Combustion

Some of New York State's municipal waste is combusted in facilities equipped to use the heat produced by the incineration process to generate electricity. This source category includes the CO₂ and N₂O emissions produced during the incineration process at these waste-to-energy facilities. Chapter 6 covers emissions associated with other forms of waste management (i.e., those that are not associated with the production of electricity).

4.2 Emissions Inventory Data and Methodology

GHG emissions related to the power supply represent emissions produced at the point of electric generation only. In addition, according to the EPA emission inventory protocols and the methodologies prescribed in the eGRID2010 Technical Support Document,⁵² the carbon dioxide emissions from the combustion of biogenic fuels (e.g., wood, landfill gas, and the biomass component of municipal waste) are not included in estimates of GHG emissions.

4.2.1 Fuel Combustion

Estimates for emissions from direct fuel used the SIT software and the methods provided in the EIIP guidance document for electric sector fuel combustion.⁵³

Electricity generation: The default historical fuel consumption data in the SIT for New York State came from the SEDS and are updated with fuel consumption data from NYSERDA's *Patterns and Trends*.⁵⁴

⁵² Environmental Protection Agency. 2014. *The Emissions & Generation Resource Integrated Database for 2007 (eGRID2007) Technical Support Document*. http://www.epa.gov/cleanenergy/documents/egridzips/eGRID_9th_edition_V1-0_year_2010_Technical_Support_Document.pdf.

⁵³ GHG emissions were calculated using SIT, with reference to EIIP. 2006 "Chapter 1: Methods for Estimating Carbon Dioxide Emissions from Combustion of Fossil Fuels.", and. "Chapter 2: Methods for Estimating Methane and Nitrous Oxide Emissions from Stationary Combustion." Volume VIII.

⁵⁴ NYSERDA. 2016. "Patterns and Trends, New York State Energy Profiles: 2000–2014" <https://www.nyserra.ny.gov/About/Publications/EA-Reports-and-Studies/Patterns-and-Trends>.

Net imports of electricity: Emissions from net imports of electricity were based on output from the Integrated Planning Model (IPM[®]), an electricity sector modeling software platform developed by ICF International, to support the development of the New York State Energy Plan. IPM is a linear programming model that incorporates electricity systems throughout the U.S. and Canada, including those managed by NYISO, the New England Independent System Operator (ISO-NE), and the Pennsylvania-New Jersey-Maryland Interconnection (PJM). Key input data include existing and planned generation units, annual electricity demand by zone, load shapes, transmission system capacities and transfer limits, generation unit level operation and maintenance costs and performance characteristics, fuel prices, new capacity and emission control technology costs and performance characteristics, zonal reliability requirements, national and state environmental regulations, and financial market assumptions. For imports, the emission factor was estimated based on modeled emissions from neighboring electric service territories.

4.2.2 Electricity Distribution

Emissions estimates from T&D used the SIT and the methods provided in the EIIP guidance document for industrial process emissions.⁵⁵ The default data assumed that SF₆ consumption is equivalent to SF₆ sales. The default historical SF₆ data in SIT came from the EIA's Electric Power Annual report.⁵⁶

4.2.3 Municipal Waste Combustion

Emissions from waste combustion were based on waste-to-energy facility-specific tonnage data from EPA's SIT module from 1990–2007 and from the New York State Department of Environmental Conservation's (NYSDEC) Solid Waste Information Management System (SWIMS) data from 2008 to 2014. These data were input into the SIT, and the methods provided in the EIIP guidance document for municipal solid waste emissions were used to estimate the CO₂ and N₂O produced.⁵⁷

⁵⁵ GHG emissions were calculated using SIT, with reference to EIIP. 2006. "Chapter 6: Methods for Estimating Non-Energy Greenhouse Gas Emissions from Industrial Processes." Volume VIII.

⁵⁶ U.S. Energy Information Administration. 2013. *Electric Power Annual 2011*. Volume I. <http://www.eia.gov/electricity/annual/archive/2011/>.

⁵⁷ GHG emissions were calculated using SIT, with reference to EIIP. "Chapter 13: Methods for Estimating Greenhouse Gas Emissions from Municipal Solid Waste." Volume VIII.

4.3 Results

Figure 4-1 provides a 2014 breakdown of electricity generation and net imports of electricity for New York by source category. The primary energy sources were natural gas (34 percent) and nuclear power (27 percent) followed by conventional hydropower (16 percent) and net imports of electricity (14 percent). Figure 4-1 shows that nearly half of the State's electricity generation comes from sources that produce little or no emissions, such as conventional hydropower, nuclear power, and wind energy.

Figure 4-1. Proportion of New York State Electricity Generation and Net Imports by Source Category, 2014

The Wood category includes wood waste.




Figure 4-2 provides a 2014 breakdown by percentage of emissions from power supply and delivery within New York State by source category. While the majority of emissions come from natural gas combustion (59 percent), coal is the second-highest source of emissions percentage-wise (10 percent), though it only provides a relatively small proportion of the State's electricity generation (2.7 percent). Net imports of

electricity also contribute approximately 19 percent of GHG emissions, however there is some uncertainty associated with the estimation of emissions from this out-of-state source category. SF₆ emissions from the transmission and distribution of electricity and petroleum emissions are the smallest sources at only one percent each of total emissions.

Figure 4-2. Proportion of GHG Emissions from Power Supply and Delivery by Source Category, 2014

CO₂ emissions from Municipal Waste Combustion are assumed to come from the non-biogenic portions of the waste only. In accordance with EPA guidelines, the biogenic portions of the waste are assumed to have no net CO₂ emissions.




Figure 4-3 shows the electricity generation and net imports of electricity for the State by source category for 1990–2014. During that time period, under existing state policies and economic conditions, the largest absolute increase in electricity generation by source comes from natural gas, which increases from approximately 23,000 GWh of generation in 1990 to approximately 54,000 in 2014. As indicated by Table 4-1, the renewables category, which includes renewable energy sources such as solar, wind, and bioenergy, saw the greatest growth percentage with an estimated average annual growth rate of 2.3 percent between 1990 and 2014. Net imports of electricity have increased substantially, from approximately 5,000 GWh in 1990 to 22,000 in 2014.

Figure 4-3. New York State Electricity Generation and Net Imports of Electricity by Source Category, 1990–2014

Note: The Other category includes wood, landfill gas, and waste. The Hydropower category includes both conventional and pumped storage hydropower.




Table 4-1. Electric Generation (GWh) and Annual Average Growth Rates by Fuel Type

The Other category includes wood, landfill gas, and waste. The Hydropower category includes both conventional and pumped storage hydropower.

Energy Source	1990	2014	Average % Annual Growth 1990 – 2014
Coal	25,913	4,325	(0.06)
Natural Gas	22,724	54,380	0.05
Petroleum	33,885	2,136	(0.01)
Hydropower	28,188	26,823	0.00
Nuclear	23,623	43,041	0.03
Net Imports of Electricity	4,519	22,103	0.15
Renewables	-	4,595	2.29
Other	-	2,655	0.02
Total	138,853	160,059	0.87

Figure 4-4 shows the primary fossil fuel energy use at in-State power stations for natural gas, coal, and petroleum from 1990–2014. The data show natural gas replacing coal and petroleum generation during this time.

Figure 4-4. Primary Fossil Fuel Energy Use at New York State Power Stations by Fuel Type




Figure 4-5 shows GHG emissions from power supply and delivery within the State by source category for 1990–2014. Between 1990 and 2014, total GHG emission from power supply and delivery (including imports) decreased by approximately 25.7 million metric tons of carbon dioxide equivalent (MMtCO₂e) to reach approximately 41.5 MMtCO₂e. Emissions of SF₆ from electrical equipment experienced declines since the mid-1990s, mostly due to voluntary action by industry. SF₆ emissions from electric power T&D were approximately 1.3 MMtCO₂e in 1990 and decreased to approximately 0.3 MMtCO₂e in 2014.

The largest absolute increase in GHG emissions between 1990 levels and 2014 status quo levels comes from natural gas, with emissions rising from approximately 12.6 MMtCO₂e in 1990 to approximately 24.7 MMtCO₂e in 2014, as indicated by Table 4-2. Emissions associated with coal and petroleum combustion decline from 24.5 MMtCO₂e in 1990 to 4.3 MMtCO₂e in 2014, and 25.9 MMtCO₂e in 1990 to 1.4 MMtCO₂e in 2014, respectively.

Figure 4-5. GHG Emissions from Power Supply and Delivery by Source Category, 1990–2014

CO₂ emissions from municipal waste combustion are assumed to come from the non-biogenic portions of the waste only. In accordance with EPA guidelines, the biogenic portions of the waste are assumed to have no net CO₂ emissions.




Table 4-2. GHG Emissions from Power Supply and Delivery by Source Category, 1990–2014 (MMtCO₂e)

CO₂ emissions from municipal waste combustion are assumed to come from the non-biogenic portions of the waste only. In accordance with EPA guidelines, the biogenic portions of the waste are assumed to have no net CO₂ emissions.

Source Category	1990	1995	2000	2005	2010	2014
Natural Gas	12.6	23.4	20.2	16.5	23.0	24.7
Coal	24.5	21.4	23.7	19.8	13.2	4.3
Petroleum	25.9	6.5	11.8	17.3	1.1	1.4
Municipal Waste Combustion	1.3	2.0	2.0	2.4	2.7	2.8
Transmission and Distribution	1.3	0.9	0.6	0.5	0.3	0.3
Net Imported Electricity	1.6	4.3	5.7	6.6	9.6	8.0
Electric Power Including Imports	67.2	58.4	63.9	63.0	49.9	41.5
Electric Power Excluding Imports	65.5	54.1	58.3	56.4	40.3	33.5

5 Agriculture

5.1 Overview

The emissions discussed in this chapter refer to non-energy methane (CH_4) and nitrous oxide (N_2O) emissions from both livestock and crop production. Energy emissions related to agricultural practices (combustion of fossil fuels to power agricultural equipment) are included in the residential, commercial, and industrial (RCI) fuel consumption sector estimates (see Chapter 1).

The inventory accounts for both direct and indirect emissions of N_2O -related to livestock and crop production. Direct emissions occur at the site of application of manure, fertilizer, and sewage sludge to agricultural soils. When nitrogen is applied to soils, indirect emissions can occur through the volatilization of ammonia and oxides of nitrogen. These products can then be re-deposited, enter the nitrification/denitrification cycle, and be emitted as N_2O in another location. Indirect emissions can also occur through leaching or runoff of nitrogen, which can enter the nitrification/denitrification cycle on or off-site and then be emitted as N_2O .

The primary agricultural GHG sources—livestock and agricultural soils—are further subdivided as described in the following sections.

5.1.1 Livestock

Enteric Fermentation: CH_4 emissions from enteric fermentation are the result of normal digestive processes in ruminant and non-ruminant livestock. Microbes in the animal digestive system break down food and emit CH_4 as a by-product. Ruminant livestock produce more CH_4 than non-ruminant livestock because of digestive activity in the large fore-stomach.

Manure management: CH_4 and N_2O emissions from the storage and treatment of livestock manure (e.g., in compost piles or anaerobic treatment lagoons) occur as a result of manure decomposition. The environmental conditions of decomposition drive the relative magnitude of emissions. In general, the more anaerobic the conditions are, the more CH_4 is produced because decomposition is aided by CH_4 -producing bacteria that thrive in oxygen-limited conditions. In contrast, N_2O emissions increase under aerobic conditions.

Emission estimates from manure management are based on manure stored and treated on livestock operations (e.g., dairies, feedlots, swine operations). Emissions from manure deposited directly on land by grazing animals and emissions from manure applied to agricultural soils as an amendment are accounted for under animal production in the following sections.

5.1.2 Agricultural Soils

Fertilizers: The application of synthetic and organic fertilizers can result in N₂O emissions. Nitrogen additions drive the underlying soil nitrification and denitrification cycle, which produces N₂O as a by-product.

Crops: This source category covers N₂O emissions from the decomposition of crop residues and the production of nitrogen fixing crops.

Animal production: This source category covers N₂O emissions resulting from animal excretions directly on agricultural soils (e.g., pasture, paddock, or range) or manure spreading on agricultural soils, including leaching and runoff.

5.2 Emissions Inventory Data and Methodology

Historical GHG emissions were estimated using the SIT software and the methods provided in the EIIP guidance document for the sector.⁵⁸ In general, the SIT methodology applies emission factors developed for the U.S. to activity data for the agriculture sector. Activity data include livestock population statistics, amounts of fertilizer applied to crops, and trends in manure management practices. This methodology is based on international guidelines developed by sector experts for preparing GHG emissions inventories.^{59,60}

⁵⁸ GHG emissions were calculated using SIT, with reference to EIIP. 2006. “Chapter 8: Methods for Estimating Greenhouse Gas Emissions from Livestock Manure Management.” and “Chapter 10: Methods for Estimating Greenhouse Gas Emissions from Agricultural Soil Management.” Volume VIII.

⁵⁹ Intergovernmental Panel on Climate Change. 1996. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*. <http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.htm>.

⁶⁰ Intergovernmental Panel on Climate Change. 2000. *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. <http://www.ipcc-nggip.iges.or.jp/public/gp/english/>.

Historical data on crop production in New York State and on the number of animals in the State were obtained from the National Agricultural Statistical Service (NASS) of the United States Department of Agriculture (USDA), and incorporated as defaults in the SIT.⁶¹ The default SIT manure management system assumptions for each livestock category were used for this inventory. SIT data on fertilizer usage came from *Commercial Fertilizers*, a report from the Fertilizer Institute and are also contained as defaults with the SIT module. Details for each of the livestock and crop production emission sources are provided below.

5.2.1 Livestock

Agricultural animals: SIT default data on livestock populations are derived from the USDA NASS. Methane emission factors specific to each type of animal by region (e.g., dairy cattle, beef cattle, sheep, goats, swine, and horses) are provided in the SIT. For years 2012–2014, New York State emissions reported in the EPA’s 2014 U.S. GHG Inventory are used.

Manure management: The same population data used for enteric fermentation are also used as input to estimate CH₄ and N₂O emissions from manure management. Population estimates are multiplied by an estimate for typical animal mass and a volatile solids (VS) production rate to estimate the total VS produced. The VS estimate for each animal type is then multiplied by a maximum potential CH₄ emissions factor and a weighted CH₄ conversion factor to derive total CH₄ emissions. The methane conversion factor adjusts the maximum potential CH₄ emissions based on the types of manure management systems employed in the State.

Nitrous oxide emissions are derived using the animal population estimates discussed above, multiplied by the typical animal mass and a total Kjeldahl nitrogen (K-nitrogen) production factor. The total K-nitrogen is multiplied by a non-volatilization factor to determine the fraction that is managed in manure management systems. The unvolatilized portion is divided into fractions that are processed in either liquid (e.g., lagoons) or solid waste management systems (e.g., storage piles, composting). Each of these fractions is then multiplied by an N₂O emission factor and the results summed to estimate total N₂O emissions. For years 2012–2014 SIT values are modified to reflect VS and K-nitrogen emissions factors used in EPA’s 2014 GHG Inventory.

⁶¹ National Agricultural Statistical Service. https://www.nass.usda.gov/Statistics_by_State/New_York/.

5.2.2 Agricultural Soils

Fertilizers: Calculations for direct and indirect emissions from fertilization use data on the amount of nitrogen applied to the soil in the form of both synthetic and organic fertilizers along with factors on the fraction of nitrogen volatilized and an IPCC-based emission factor for N₂O emissions from the re-deposited nitrogen (0.01 kg N₂O-N/kg N re-deposited).

Crops: Default data on tonnage of crop production from the USDA NASS were used to calculate N₂O emissions from crop residues and crops that fix nitrogen.

Animal production: The calculation of emissions for this source category requires data on animal population, mass and nitrogen emitted per unit of animal mass, as well as the amount of manure left on the soil.

To calculate indirect emissions from this source category, data on nitrogen inputs from animals to crop soils are used along with factors on the fraction of nitrogen volatilized and an IPCC-based emission factor for N₂O emissions from the re-deposited nitrogen (0.01 kg N₂O-N/kg N re-deposited).

5.3 Results

Annual GHG emissions from agricultural sources in the State range between approximately 5.3 and 6.3 MMtCO₂e during the time period between 1990 and 2014, generally staying stable over time, as shown in Figure 5-1. Detailed information on GHG emissions from agricultural sources is provided in Table 5-1. Enteric fermentation accounted for about 58 percent (3.4 MMtCO₂e) of total agricultural emissions in both 1990 and about 57 percent (3.6 MMtCO₂e) in 2014. The manure management category accounted for 14 percent (0.8 MMtCO₂e) of total agricultural emissions in 1990 and accounts for an estimated 19 percent (1.2 MMtCO₂e) of total agricultural emissions in 2014. The agricultural soils category shows 1990 emissions accounting for 28 percent (1.6 MMtCO₂e) of total agricultural emissions and 2014 emissions estimated to be about 25 percent (1.6 MMtCO₂e) of total agricultural emissions.

Figure 5-1. GHG Emissions from Agriculture, 1990–2014

The Agricultural Soil Management category includes emissions from fertilizers, crops (crop residues and nitrogen fixing crops), and animal production. Soil carbon sequestration is not shown (see Table 5-1).




Table 5-1. GHG Emissions from Agriculture, 1990–2014 (MMtCO₂e)

Source Category	1990	1995	2000	2005	2010	2014
Enteric Fermentation	3.4	3.2	3.3	3.2	3.4	3.6
Manure Management	0.8	0.8	1.0	1.1	1.1	1.2
Agriculture Soil Management	1.6	1.6	1.6	1.6	1.6	1.6
Total	5.8	5.6	5.8	5.8	6.0	6.3

6 Waste Management

6.1 Overview

The sources of GHG emissions from waste management included in this inventory report cover both solid waste and wastewater. These primary sources are further subdivided as follows:

6.1.1 Landfills

Municipal solid waste landfills: Methane (CH_4) emissions are generated from the anaerobic decomposition of the organic matter present in landfilled waste by methanogenic bacteria. Some municipal solid waste (MSW) landfills employ control technologies, such as flares that convert the CH_4 portion of recovered landfill gas to carbon dioxide (CO_2), to reduce the emissions of GHGs to the atmosphere.

Industrial solid waste landfills: This source category covers CH_4 emissions produced from waste discarded in non-hazardous industrial landfills.

6.1.2 Municipal Wastewater Management

This source category covers both CH_4 and N_2O emissions produced at municipal wastewater treatment facilities.

In addition to the sources outlined in previous sections, emissions can also result from industrial wastewater treatment, commercial/industrial waste incineration, and MSW combustion for energy production (waste-to-energy combustion). Only municipal wastewater treatment sources were considered based on the information presently available at the State level. There is no significant commercial/industrial waste incineration within the State, and waste-to-energy combustion emissions are accounted for in Chapter 4. Note that this inventory does not currently capture any other forms of waste combustion, including medical waste or hazardous waste incineration.

6.2 Emissions Inventory Data and Methodology

6.2.1 Landfills

Municipal solid waste landfills: For municipal solid waste landfills, emissions for 1990–2014 were calculated in the SIT using tonnage data from EPA’s SIT module from 1990–2007 and the DEC’s 2008–2014 SWIMS data, as well as estimates of methane flaring and capture from the EPA Landfill Methane Outreach Program (LMOP) database.⁶²

6.2.2 Wastewater Management

For municipal wastewater treatment, emissions calculations used the SIT methodology, which is based on State population, assumed biochemical oxygen demand (BOD) and protein consumption per capita, and emission factors for N₂O and CH₄. Table 6-1 shows the key SIT default values. Municipal wastewater treatment emissions were, therefore, based on the population growth rate for 1990–2014.

Table 6-1. SIT Key Default Values for Municipal Wastewater Treatment

Source: U.S. EPA State Inventory Tool – Wastewater Module; methodology and factors taken from U.S. EPA, Emission Inventory Improvement Program, Volume 8, Chapter 12, December 2006.

Variable	Default Value
BOD	0.09 kg /day-person
Amount of BOD anaerobically treated	16.25%
CH ₄ emission factor	0.6 kg/kg BOD
New York residents not on septic	79%
Water treatment N ₂ O emission factor	4.0 g N ₂ O/person-yr
Biosolids emission factor	0.005 kg N ₂ O-N/kg sewage-N

⁶² Environmental Protection Agency. 2016. “Landfill Methane Outreach Project Index.” Last modified October 4, 2016. <http://www.epa.gov/lmop/projects-candidates/index.html>.

6.3 Results

The emission estimates for the waste management sector, shown in Figure 6-1 and Table 6-2, include GHG emissions associated with waste generated in the State. Overall, the sector accounted for 14.8 MMtCO₂e in 1990 and 13.8 MMtCO₂e in 2014. The share of emissions from landfills decreases slightly over the analysis period from 87 percent of the sector total in 1990 to 85 percent in 2014, while the share of emissions from wastewater increases from 13 percent in 1990 to 15 percent in 2014.

Figure 6-1. GHG Emissions from Waste Management, 1990–2014




Table 6-2. GHG Emissions from Waste Management (MMtCO₂e), 1990–2014

Source Category	1990	1995	2000	2005	2010	2012	2014
Landfills	12.89	13.43	13.52	13.49	12.17	12.06	11.63
Municipal Wastewater	1.91	1.94	2.04	2.07	2.09	2.11	2.12
Total	14.81	15.37	15.56	15.56	14.26	14.17	13.75

Many of the materials that comprise MSW – aluminum cans, steel cans, glass, plastic, paper, etc. – represent significant embedded GHG emissions, which are also not included. Embedded emissions include emissions from the entire energy-cycle of a material, including emissions from raw material extraction, transportation of the raw material, manufacture of a product or packaging, and transportation of the good or packaging to the marketplace.

Appendix A

Table A-1. Fuel Combustion Emission Factors by Sector

Values represent aggregate CO₂, CH₄ and N₂O emissions

Fuel Type	Emission Factor (lb CO₂e / MMBtu)
Aviation Fuel	159.2
Coal	204.9
Distillate Fuel Oil (No. 2)	162.9
Gasoline	158.0
Kerosene	161.2
Natural Gas	117.2
Propane/Liquefied Petroleum Gas	136.1
Residual Fuel Oil (No. 6)	166.0
Wood	18.2
Source: Emissions Factors were derived from the U.S. EPA's State Inventory Tool (release Feb, 2016)	

A.1 Global Warming Potentials

Greenhouse Gas	Global Warming Potential
CO ₂	1
CH ₄	25
N ₂ O	298
HFC-23	14,800
HFC-32	675
HFC-125	3,500
HFC-134a	1,430
HFC-143a	4,470
HFC-152a	124
HFC-227ea	3,220
HFC-236fa	9,810
HFC-4310mee	1,640
PFCs	7,390-13,300
SF ₆	22,800
Source: IPCC Fourth Assessment Report (2007)	

NYSERDA, a public benefit corporation, offers objective information and analysis, innovative programs, technical expertise, and support to help New Yorkers increase energy efficiency, save money, use renewable energy, and reduce reliance on fossil fuels. NYSERDA professionals work to protect the environment and create clean-energy jobs. NYSERDA has been developing partnerships to advance innovative energy solutions in New York State since 1975.

To learn more about NYSERDA's programs and funding opportunities, visit nyserda.ny.gov or follow us on Twitter, Facebook, YouTube, or Instagram.

**New York State
Energy Research and
Development Authority**

17 Columbia Circle
Albany, NY 12203-6399

toll free: 866-NYSERDA
local: 518-862-1090
fax: 518-862-1091

info@nyserda.ny.gov
nyserda.ny.gov



NYSERDA

State of New York

Andrew M. Cuomo, Governor

New York State Energy Research and Development Authority

Richard L. Kauffman, Chair | John B. Rhodes, President and CEO

